



## Caringbah High School

# 2015

## Trial HSC Examination

# Mathematics Extension 2

### General Instructions

- Reading time – 5 minutes
- Working time – 3 hours
- Write using black or blue pen (Black pen is preferred)
- Board-approved calculators may be used
- A table of standard integrals is provided at the back of this paper
- In Questions 11–16, show relevant mathematical reasoning and/or calculations

### Total marks – 100

#### **Section I** Pages 2 – 5

#### **10 marks**

- Attempt Questions 1–10
- Allow about 15 minutes for this section

#### **Section II** Pages 6 – 14

#### **90 marks**

- Attempt Questions 11–16
- Allow about 2 hour and 45 minutes for this section

**Question 1 - 10** (1 mark each) Answer on page provided.

**1** The argument of  $z = -3i$  is:

- A)  $\frac{\pi}{2}$                       B)  $\frac{3\pi}{2}$                       C)  $-\frac{\pi}{2}$                       D)  $-\frac{3\pi}{2}$

**2** Consider a polynomial  $P(x)$  that has real coefficients. Which of the following could not be a pair of possible solutions to  $P(x) = 0$ ?

- A)  $x_1 = -1 + i; x_2 = -1 - i$                       B)  $x_1 = 1 + i; x_2 = 1 - i$   
 C)  $x_1 = -2 + i; x_2 = 2 - i$                       D)  $x_1 = -2 + i; x_2 = -2 - i$


**3** Given that  $z^3 = 1$  and  $\omega$  is a complex solution, the value of  $\omega^3 + \omega^4 + \omega^5$  is:

- A) 0    B) 1  
 C)  $\omega$     D)  $-\omega$

**4** The coordinates of the foci of the hyperbola  $\frac{x^2}{4} - y^2 = 1$  are given by:

- A)  $(0, \pm\sqrt{5})$     B)  $(\pm\sqrt{5}, 0)$   
 C)  $(0, \pm 2\sqrt{5})$     D)  $(\pm 2\sqrt{5}, 0)$

- 5 Which of the following is the equation of the circle shown below?



- A)  $(z + 2)(\bar{z} + 2) = 4$                       B)  $(z - 2)(\bar{z} - 2) = 4$
- C)  $(z + 2i)(\bar{z} - 2i) = 4$                       D)  $(z - 2i)(\bar{z} + 2i) = 4$

- 6 The roots of the polynomial  $4x^3 + 4x - 5 = 0$  are  $\alpha, \beta$  and  $\gamma$ .

What is the value of  $(\beta + \gamma - 3\alpha)(\alpha + \beta - 3\gamma)(\gamma + \alpha - 3\beta)$ ?

- A) 16    B) 80
- C) -16    D) -80

- 7 The equation  $x^2 + 2y^2 - 2xy + x = 8$  defines  $y$  implicitly as a function of  $x$ .

What is the value of  $\frac{dy}{dx}$  at the point  $(3, 2)$ ?

- A)  $\frac{1}{4}$     B)  $-\frac{1}{4}$
- C)  $\frac{3}{2}$     D)  $-\frac{3}{2}$

- 8 All of the integrals below are of the form  $\int_{-1}^1 f(x) dx$ .

Without evaluating, which of the integrals can be rewritten as  $2\int_0^1 f(x) dx$ ?

A)  $\int_{-1}^1 e^x \tan^{-1}(x^2) dx$

B)  $\int_{-1}^1 \frac{x^2 \sin x}{x^2 + 5} dx$

C)  $\int_{-1}^1 \sqrt{x^2 + e^x} dx$

D)  $\int_{-1}^1 x^3 \sin^{-1} x dx$

- 9 Using the recurrence relation  $U_n = \int \tan^n x dx = \frac{\tan^{n-1} x}{n-1} - U_{n-2}$

what is the value of  $\int \tan^6 x dx$ ?


A)  $\frac{\tan^5 x}{5} - \frac{\tan^4 x}{4} + \frac{\tan^3 x}{3} - \frac{\tan^2 x}{2} + \tan x + c$

B)  $\frac{\tan^5 x}{5} - \frac{\tan^3 x}{3} + \tan x + c$

C)  $\frac{\tan^5 x}{5} - \frac{\tan^3 x}{3} + \tan x - x + c$

D)  $\frac{\tan^5 x}{5} - \frac{\tan^3 x}{3} - \tan x - x + c$

10



The region bounded by the  $x$ -axis, the curve  $y = \sqrt{x^2 - 1}$  and the line  $x = 2$  is rotated around the  $y$ -axis.

The slice at  $P(x, y)$  on the curve is perpendicular to the axis of rotation.

What is the volume  $\delta V$  of the annular slice formed?

- A)  $\pi(3 - y^2)\delta y$                       B)  $\pi(4 - (y^2 + 1)^2)\delta y$
- C)  $\pi(4 - x^2)\delta x$                       D)  $\pi(2 - x^2)\delta x$

**END OF MULTIPLE CHOICE QUESTIONS**

**Section II****60 marks****Attempt all questions 11–14****Allow about 1 hour and 45 minutes for this section**

Answer each question in a SEPARATE writing booklet. Extra writing booklets are available.


In Questions 11–14, your responses should include relevant mathematical reasoning and/or calculations.

- | <b>Question 11</b> (15 marks) Start a NEW booklet.   | <b>Marks</b> |
|--|--------------|
| a) Let $z = 1 + i$ and $u = 2 - i$ . Find:   |              |
| (i) $\text{Im}(uz)$ .  | 1            |
| (ii) $ u - z $ .   | 1            |
| (iii) $-i\bar{u}$ .  | 1            |
| b) Evaluate $\int_0^1 x^2 \sqrt{1-x} \, dx$ .  | 3            |
| c) It is given that $Z = \frac{\sqrt{6}}{2} + \frac{\sqrt{2}}{2}i$ .   |              |
| (i) Show that $Z$ can be expressed in the form $\sqrt{2} \text{cis} \frac{\pi}{6}$ .   | 2            |
| (ii) Hence, express $Z^8$ in $x + iy$ form.  | 2            |
| d) (i) Prove that if the polynomial $P(x)$ has a root $\alpha$ of multiplicity $m$<br>then $P'(x)$ has a root $\alpha$ of multiplicity $m - 1$ . | 2            |
| (ii) Hence, given that the polynomial $P(x) = x^4 + x^3 - 3x^2 - 5x - 2$ has<br>a root of multiplicity 3, find all the roots of $P(x)$ .         | 3            |

**Question 12** (15 marks) Start a NEW booklet.

**Marks**

a) Consider the function  $y = f(x)$  drawn below.



On separate sketches, showing any important features, neatly draw the graphs of

- |       |  |   |
|-------|--|---|
| (i)   | $y = 1 - f(x)$                                     | 1 |
| (ii)  | $y = \sqrt{f(x)}$                                  | 1 |
| (iii) | $y = f\left(\frac{1}{x}\right)$                    | 2 |
| (iv)  | $y = e^{f(x)}$                                     | 2 |
| b)    | Evaluate $\int_8^{12} \frac{1}{x^2 - 16x + 80} dx$ | 3 |

**Question 12 continues on page 8**

Question 12 (continued)

Marks

c) The equation  $x^3 + 2x - 1 = 0$  has roots  $\alpha$ ,  $\beta$  and  $\gamma$ . Find

(i) the value of  $\alpha^2 + \beta^2 + \gamma^2$ . 2

(ii) the value of  $\alpha^3 + \beta^3 + \gamma^3$ . 2

(iii) the equation with roots  $-\alpha$ ,  $-\beta$ ,  $-\gamma$ . 2

**End of Question 12**



**Question 13** (15 marks) Start a NEW booklet.**Marks**

- a) Consider the locus of the point  $P(x, y)$  whose co-ordinates satisfy the parametric equations  $x = \cos t$ ,  $y = 1 - \cos 2t$ .
- (i) State the domain and range of the locus of  $P$ . 2
- (ii) Determine the Cartesian equation of the locus of  $P$ . 2
- (iii) Neatly sketch the curve traced by the point  $P(x, y)$ . 1
- b) Find  $\int \frac{\cos x}{\sin x + \sin^2 x} dx$ . 4
- c) An ellipse has the equation  $\frac{x^2}{16} + \frac{y^2}{9} = 1$
- (i) Prove that the tangent to the ellipse at  $P(4\cos\theta, 3\sin\theta)$  has 3  
equation  $\frac{x \cos \theta}{4} + \frac{y \sin \theta}{3} = 1$ .
- (ii) The ellipse meets the  $y$ -axis at  $B$  and  $B'$ . The tangents at  $B$  and  $B'$  3  
meet the tangent at  $P$  at the points  $Q$  and  $Q'$ .  
Find  $BQ \times B'Q'$ .

**Question 14** (15 marks) Start a NEW booklet.**Marks**


- a) Consider the functions  $f(x) = |x| + 1$  and  $g(x) = \frac{6}{|x|}$ .
- (i) Solve the equation  $f(x) = g(x)$ . 2
- (ii) Sketch the graphs of  $y = f(x)$  and  $y = g(x)$  on the same axes. 2
- (iii) Hence or otherwise solve the inequality  $g(x) > f(x)$ . 1
- b) i) Prove that  $a^2 + b^2 \geq 2ab$ . 1
- ii) Hence or otherwise prove that  $(p + 2)(q + 2)(p + q) \geq 16pq$  2  
where  $p$  and  $q$  are positive real numbers.
- c) The base of a solid is the region between the lines  $y = 3x$  and  $y = -x$  from  $x = 0$  to  $x = 2$ .  
Each cross section by planes perpendicular to the  $x$ -axis is a square.  
Calculate the volume of the solid. 3

**Question 14 continues on page 11**

## Question 14 (continued)

Marks

- d) In the diagram below the two circles touch internally at  $A$ .  $O$  is the centre of the larger circle.  $B$  is a point on the larger circle and chord  $AB$  cuts the smaller circle at  $S$ .



**Answer this question on the page provided**

Prove that chord  $AB$  is bisected at  $S$ .

4

[ Ensure that any constructions are clearly shown and labelled ]

**End of Question 14**

**Question 15** (15 marks) Start a NEW booklet.

**Marks**

a) Let  $z$  be a complex number such that  $|z| = r$  and  $\arg z = \theta$  for  $0 < \theta < \frac{\pi}{2}$ .

Prove that  $\arg(r^2 - z^2) = \theta - \frac{\pi}{2}$ .

3

b) (i) Show that  $\frac{d}{dx}\left(\frac{x^2}{2}\ln x - \frac{x^2}{4}\right) = x \ln x$

1

(ii) Using  $\frac{1}{3}$  of a page neatly sketch the region in the number plane that contains all points satisfying simultaneously the inequalities

$$x \leq 1, y \geq 1, y \leq e^x.$$

1

(iii) This region is rotated through one complete revolution about the  $x$ -axis.

Use the method of cylindrical shells to find the volume of the resulting solid.

4

c) If  $n$  is a positive integer and  $f(x) = e^{-x}\left(1 + x + \frac{x^2}{2!} + \dots + \frac{x^n}{n!}\right)$ ,  $x \geq 0$

(i) Show that  $f(x)$  is a decreasing function.

3

(ii) Deduce that for  $x > 0$  and  $n$  any positive integer,

3

$$e^x \geq 1 + x + \frac{x^2}{2!} + \dots + \frac{x^n}{n!}$$


**Question 16** (15 marks) Start a NEW booklet.

**Marks**

a) Find  $\int \frac{e^{2x}}{e^x - 1} dx$ .

3

b)



The point  $P\left(cp, \frac{c}{p}\right)$  lies on the hyperbola  $xy = c^2$ . The point  $T$  lies at the foot of the perpendicular drawn from the origin  $O$  to the tangent at  $P$ .

(i) Show that the tangent at  $P$  has equation  $x + p^2y = 2cp$ . 2

(ii) If the coordinates of  $T$  are  $(x_1, y_1)$  show that  $y_1 = p^2x_1$ . 1

(iii) Show that the locus of  $T$  is given by  $(x^2 + y^2)^2 = 4c^2xy$ . 2

**Question 16 continues on page 14**

Question 16 (continued)

Marks

c) Consider the integral  $I_n = \int_0^1 x^{2n+1} e^{-x^2} dx$ .

(i) Use integration by parts to show that  $I_n = -\frac{1}{2e} + nI_{n-1}$ , for  $n \geq 1$ . 2

(ii) Show that  $I_0 = \frac{1}{2} - \frac{1}{2e}$  and  $I_1 = \frac{1}{2} - \frac{1}{e}$  1

(iii) Prove by mathematical induction that for all  $n \geq 1$ : 3

$$1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \dots + \frac{1}{n!} = e - \frac{2eI_n}{n!}$$

(iv) It is given that  $0 \leq I_n \leq 1$  because  $0 \leq x^{2n+1} e^{-x^2} \leq 1$ , for  $0 \leq x \leq 1$ .


**[Do not prove this]**

Use this fact to help to evaluate  $\frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \dots$  1

giving your answer in exact form.

**END OF EXAM**

Question 14d (additional diagram) Candidate Name/Number: \_\_\_\_\_



**Hand in this page inside the Question 14 booklet**

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

---

**Multiple Choice Section:**

- 1.C      2.C      3.A      4.B      5.B  
 6.D      7.D      8.D      9.C      10.A

**Question 1.**

$-\frac{\pi}{2}$  ----- [C]

**Question 2.**

Solutions need to be complex conjugates as coefficients are real

$x_1 = -2 + i; x_2 = 2 - i$  are not ----- [C]

**Question 3.**

$\omega^3 - 1 = 0 \rightarrow (\omega - 1)(1 + \omega + \omega^2) = 0$   
 and since  $\omega \neq 1$  ( must be non-real ), then  
 $\omega^3 + \omega^4 + \omega^5 = 1 + \omega + \omega^2$   
 $= 0$  ----- [A]

**Question 4.**

$a = 2; b = 1; b^2 = a^2(e^2 - 1); \text{Foci } (\pm ae, 0)$   
 $\therefore e^2 = \frac{5}{4} \rightarrow e = \frac{\sqrt{5}}{2}$   
 $\therefore \text{Foci } (\pm\sqrt{5}, 0)$  ----- [B]

**Question 5.**

Equation in Cartesian form is  $(x - 2)^2 + y^2 = 4$   
 $(z - 2)(\bar{z} - 2) = 4 \rightarrow z\bar{z} - 2(z + \bar{z}) + 4 = 4$   
 $\therefore x^2 + y^2 - 2(2x) + 4 = 4$   
 $\therefore x^2 - 4x + 4 + y^2 = 4$   
 $\therefore (x - 2)^2 + y^2 = 4$  ----- [B]

**Question 6.**

$(\beta + \gamma - 3\alpha)(\alpha + \beta - 3\gamma)(\gamma + \alpha - 3\beta)$   
 $= (\alpha + \beta + \gamma - 4\alpha)(\alpha + \beta + \gamma - 4\beta)(\alpha + \beta + \gamma - 4\gamma)$   
 Now  $\alpha + \beta + \gamma = -\frac{b}{a} = 0$  and  $\alpha\beta\gamma = \frac{c}{a} = \frac{5}{4}$   
 $\therefore = (-4\alpha)(-4\beta)(-4\gamma)$   
 $= -64\alpha\beta\gamma$   
 $= -64 \times \frac{5}{4} = -80$  ----- [D]

**Question 7.**

$x^2 + 2y^2 - 2xy + x = 8$   
 $\therefore 2x + 4y \frac{dy}{dx} - 2\left(x \times \frac{dy}{dx} + y \times 1\right) + 1 = 0$   
 $\therefore 2(2y - x) \frac{dy}{dx} = 2y - 2x - 1$   
 $\therefore \frac{dy}{dx} = \frac{2y - 2x - 1}{2(2y - x)}$  and at  $P(3, 2)$   
 $\frac{dy}{dx} = \frac{4 - 6 - 1}{2(4 - 3)} = -\frac{3}{2}$  ----- [D]

**Question 8.**

Let  $f(x) = x^3 \sin^{-1} x$   
 $\therefore f(-x) = (-x)^3 \sin^{-1}(-x)$   
 $= -x^3 \times -\sin^{-1} x$   
 $= x^3 \sin^{-1} x = f(x)$   
 hence  $f(x)$  is an even function  
 hence ----- [D]



**Question 9.**

$$U_6 = \frac{\tan^5 x}{5} - U_4$$


$$U_4 = \frac{\tan^3 x}{3} - U_2$$

$$U_2 = \frac{\tan x}{1} - U_0$$

$$U_0 = \int 1 dx = x + c$$

$$\therefore U_6 = \frac{\tan^5 x}{5} - \frac{\tan^3 x}{3} + \tan x - x + c \quad \text{--- [C]}$$

**Question 10.**



where  $R = 2$   
and  $r = x$

$$\begin{aligned} \therefore \text{Area of slice} &= \pi(2^2 - x^2) \\ &= \pi(4 - (y^2 + 1)) \end{aligned}$$

$$\therefore \text{Volume of slice} = \pi(3 - y^2) \delta y \quad \text{----- [A]}$$

**Question 11**

a) i)  $\text{Im}[(1+i)(2-i)] = \text{Im}(3+1i) = 1$

ii)  $|u-z| = |1-2i| = \sqrt{5}$

iii)  $-i\bar{u} = -i \times (2+i) = 1-2i$

b) Let  $u=1-x \rightarrow x=1-u$  and  $du=-dx$   
When  $x=0, u=1; x=1, u=0$ .

$$\begin{aligned} \therefore I &= -\int_1^0 (1-u)^2 \sqrt{u} du \\ &= \int_0^1 u^{\frac{1}{2}} - 2u^{\frac{3}{2}} + u^{\frac{5}{2}} du \\ &= \left[ \frac{2}{3}u^{\frac{3}{2}} - \frac{4}{5}u^{\frac{5}{2}} + \frac{2}{7}u^{\frac{7}{2}} \right]_0^1 \\ &= \frac{2}{3} - \frac{4}{5} + \frac{2}{7} = \frac{16}{105} \end{aligned}$$

c) i)  $|Z| = r = \sqrt{\frac{6}{4} + \frac{2}{4}} = \sqrt{2}$

$$\arg Z = \theta = \tan^{-1} \left( \frac{\sqrt{2}/2}{\sqrt{6}/2} \right)$$

$$\therefore \theta = \tan^{-1} \left( \frac{1}{\sqrt{3}} \right) = \frac{\pi}{6}$$

ii)  $Z^8 = \left( \sqrt{2} \text{cis} \left( \frac{\pi}{6} \right) \right)^8$   
 $= 2^4 \text{cis} \left( \frac{8\pi}{6} \right) \quad \text{DMT}$

$$= 16 \left( \cos \frac{4\pi}{3} + i \sin \frac{4\pi}{3} \right)$$

$$= 16 \left( -\frac{1}{2} - \frac{\sqrt{3}}{2}i \right) = -8(1 + \sqrt{3}i)$$

d) i) Let  $P(x) = (x-\alpha)^m Q(x)$

$$\begin{aligned} \therefore P'(x) &= m(x-\alpha)^{m-1} Q(x) + (x-\alpha)^m Q'(x) \\ &= (x-\alpha)^{m-1} (mQ(x) + (x-\alpha)Q'(x)) \end{aligned}$$

hence  $P'(x)$  has a root  $\alpha$  of multiplicity  $m-1$ .

ii)  $P(x) = x^4 + x^3 - 3x^2 - 5x - 2$

$$P'(x) = 4x^3 + 3x^2 - 6x - 5$$

$$P''(x) = 12x^2 + 6x - 6$$

$$P''(x) = 0 \rightarrow 2x^2 + x - 1 = 0$$

$$\therefore (2x-1)(x+1) = 0 \rightarrow x = \frac{1}{2} \text{ or } x = -1$$

when  $x = -1, P'(-1) = -4 + 3 + 6 - 5 = 0$

when  $x = -1, P(-1) = 1 - 1 - 3 + 5 - 2 = 0$


hence  $x = -1$  is a root of multiplicity 3

$\therefore P(x) = (x+1)^3(x-2)$  by inspection


hence  $P(x)$  has roots  $x = -1, -1, -1, 2$ .

**Question 12.**


a) i)




ii)



iii)



iv)



$$\begin{aligned}
 \text{b) } I &= \int_8^{12} \frac{1}{(x^2 - 16x + 64) + 16} dx \\
 &= \int_8^{12} \frac{1}{(x-8)^2 + 4^2} dx \\
 &= \frac{1}{4} \left[ \tan^{-1} \left( \frac{x-8}{4} \right) \right]_8^{12} \\
 &= \frac{1}{4} (\tan^{-1} 1 - \tan^{-1} 0) = \frac{\pi}{16}
 \end{aligned}$$

c) i)

$$\begin{aligned}
 \alpha^2 + \beta^2 + \gamma^2 &= (\alpha + \beta + \gamma)^2 - 2(\alpha\beta + \beta\gamma + \gamma\alpha) \\
 &= (0)^2 - 2(2) = -4
 \end{aligned}$$

$$\text{ii) } x^3 + 2x - 1 = 0 \rightarrow x^3 = 1 - 2x$$

$$\therefore \alpha^3 = 1 - 2\alpha$$

$$\beta^3 = 1 - 2\beta$$

$$\gamma^3 = 1 - 2\gamma$$

$$\begin{aligned}
 \therefore \alpha^3 + \beta^3 + \gamma^3 &= 3 - 2(\alpha + \beta + \gamma) \\
 &= 3 - 2(0) = 3
 \end{aligned}$$

$$\text{iii) Let } x = -\alpha \rightarrow \alpha = -x$$

$\therefore$  the required equation is given by

$$\therefore (-x)^3 + 2(-x) - 1 = 0$$

$$\therefore -x^3 - 2x - 1 = 0$$

$$\therefore x^3 + 2x + 1 = 0$$

**Question 13.**


$$\text{a) i) D: } -1 \leq x \leq 1; \text{ R: } 0 \leq y \leq 2$$

$$\text{ii) } y = 1 - (2\cos^2 t - 1)$$

$$= 2 - 2(\cos t)^2$$

$$\therefore y = 2 - 2x^2$$

iii)



$$13b) I = \int \frac{\cos x}{\sin x + \sin^2 x} dx$$

$$\text{Let } u = \sin x \rightarrow du = \cos x dx$$

$$\therefore I = \int \frac{1}{u+u^2} du = \int \frac{1}{u(1+u)} du$$

$$\text{Now } \frac{1}{u(1+u)} = \frac{1}{u} - \frac{1}{u+1} \text{ using partial fractions}$$

$$\therefore I = \int \frac{1}{u} - \frac{1}{u+1} du$$

$$= \ln u - \ln(1+u)$$

$$= \ln\left(\frac{u}{1+u}\right)$$

$$\therefore I = \ln\left(\frac{\sin x}{1+\sin x}\right) + c$$

$$c) i) \frac{x^2}{16} + \frac{y^2}{9} = 1 \rightarrow a = 4, b = 3$$

$$\therefore \frac{2x}{16} + \frac{2y}{9} \cdot \frac{dy}{dx} = 0 \rightarrow \frac{dy}{dx} = -\frac{9x}{16y}$$

$$\therefore \text{ at } P(4\cos\theta, 3\sin\theta), \frac{dy}{dx} = -\frac{36\cos\theta}{48\sin\theta} = -\frac{3\cos\theta}{4\sin\theta}$$


$$\therefore \text{ eq}^n \text{ of } T \text{ at } P: y - 3\sin\theta = -\frac{3\cos\theta}{4\sin\theta}(x - 4\cos\theta)$$

$$4y \sin\theta - 12\sin^2\theta = -3x \cos\theta + 12\cos^2\theta$$

$$3x \cos\theta + 4y \sin\theta = 12(\sin^2\theta + \cos^2\theta)$$

$$\therefore \frac{x \cos\theta}{4} + \frac{y \sin\theta}{3} = 1 \left[ \sin^2\theta + \cos^2\theta = 1 \right]$$

13c)ii)



$$\text{At } Q: y = 3 \rightarrow x = \frac{4(1 - \sin\theta)}{\cos\theta}$$

$$\text{At } Q': y = -3 \rightarrow x = \frac{4(1 + \sin\theta)}{\cos\theta}$$

$$\begin{aligned} \therefore BQ \times B'Q' &= \frac{4(1 - \sin\theta)}{\cos\theta} \times \frac{4(1 + \sin\theta)}{\cos\theta} \\ &= \frac{16(1 - \sin^2\theta)}{\cos^2\theta} = 16 \end{aligned}$$

#### Question 14.

$$a) i) |x| + 1 = \frac{6}{|x|}$$


$$\therefore |x|^2 + |x| - 6 = 0$$

$$\therefore (|x| + 3)(|x| - 2) = 0$$

$$\therefore |x| = -3 \rightarrow \text{no solution}$$

$$\text{or } |x| = 2 \rightarrow x = \pm 2$$

ii)



iii)  $-2 < x < 0$  and  $0 < x < 2$

b) i)  $(a-b)^2 \geq 0$   
 $\therefore a^2 - 2ab + b^2 \geq 0$   
 $\therefore a^2 + b^2 \geq 2ab$


ii) using (i) let  $a = \sqrt{p}$  and  $b = \sqrt{q}$   
 $\therefore \sqrt{p^2} + \sqrt{q^2} \geq 2\sqrt{p}\sqrt{q}$   
 $\therefore p + q \geq 2\sqrt{p}\sqrt{q}$  ----- [1]

similarly let  $a = \sqrt{p}$  and  $b = \sqrt{2}$   
 $\therefore p + 2 \geq 2\sqrt{p}\sqrt{2}$  ----- [2]

also let  $a = \sqrt{q}$  and  $b = \sqrt{2}$   
 $\therefore q + 2 \geq 2\sqrt{q}\sqrt{2}$  ----- [3]

hence [1]  $\times$  [2]  $\times$  [3] gives:

$(p+2)(q+2)(p+q) \geq 2\sqrt{p}\sqrt{q} \times 2\sqrt{p}\sqrt{2} \times 2\sqrt{q}\sqrt{2}$   
 $\therefore (p+2)(q+2)(p+q) \geq 16(\sqrt{p})^2(\sqrt{q})^2$   
 $\therefore (p+2)(q+2)(p+q) \geq 16pq$




The area of a typical cross section is  $\delta A = (y_1 + y_2)^2$

$\therefore \delta A = (3x + x)^2 = 16x^2$

$\therefore \delta V = 16x^2 \delta x$

$\therefore V = 16 \int_0^2 x^2 dx$

$= 16 \left[ \frac{x^3}{3} \right]_0^2 = \frac{128}{3} u^3$



Construct tangent  $TA$  at  $A$ . Construct diameter  $AOC$ .  
 Join  $OS$  and  $BC$ .

Let  $\angle TAS = \alpha$ .

$\therefore \angle SOA = \alpha$  [ $\angle$  in alternate segment ( $\Delta AOS$ )]


$\therefore \angle BCA = \alpha$  [ $\angle$  in alternate segment ( $\Delta BCA$ )]

$\therefore \angle SOA = \angle BCA$  and since they are equal corresponding angles then  $BC$  is parallel to  $OS$ .

Also  $OA = OC$  (equal radii)  
 and since a family of parallel lines divides transversals in the same ratio, then  $BS = SA$ .

Hence chord  $AB$  is bisected at  $S$ .  
 [Note: Congruent triangles can also be used.]

**Question 15.**



$\arg(r^2 - z^2) = \arg(r - z) + \arg(r + z)$

Now  $\arg(r + z) = \frac{\theta}{2}$  (by isosceles  $\Delta$ )


also  $\arg(r - z) = \frac{\theta}{2} - \frac{\pi}{2}$  (by isosceles  $\Delta$ )

$\therefore \arg(r^2 - z^2) = \frac{\theta}{2} - \frac{\pi}{2} + \frac{\theta}{2} = \theta - \frac{\pi}{2}$

15b) i)

$$\begin{aligned} \frac{d}{dx} \left( \frac{x^2}{2} \ln x - \frac{x^2}{4} \right) &= \frac{x^2}{2} \times \frac{1}{x} + \frac{2x}{2} \times \ln x - \frac{2x}{4} \\ &= \frac{x}{2} + x \times \ln x - \frac{x}{2} = x \ln x \end{aligned}$$

ii)



Taking the middle of the shell at a distance 'y' units from the x-axis gives the following two radii:

$$R = y + \frac{\delta y}{2} \text{ and } r = y - \frac{\delta y}{2}$$

$\therefore$  Volume  $\delta V$  of typical cylinder =  $\pi(R+r)(R-r)h$

$$= \pi \left( y + \frac{\delta y}{2} + y - \frac{\delta y}{2} \right) \left( y + \frac{\delta y}{2} - y + \frac{\delta y}{2} \right) h$$

$$= 2\pi y h \delta y \text{ where } h = 1 - x$$

$$\text{Now } y = e^x \rightarrow x = \ln y \text{ [when } x=1, y=e \text{]}$$

$$\therefore V = 2\pi \int_1^e y(1 - \ln y) dy$$

$$\therefore V = 2\pi \int_1^e y dy - 2\pi \int_1^e y \ln y dy$$

$$= 2\pi \left[ \frac{y^2}{2} \right]_1^e - 2\pi \left[ \frac{y^2}{2} \ln y - \frac{y^2}{4} \right]_1^e$$

$$= 2\pi \left( \frac{e^2}{2} - \frac{1}{2} \right) - 2\pi \left[ \left( \frac{e^2}{2} \ln e - \frac{e^2}{4} \right) - \left( \frac{1}{2} \ln 1 - \frac{1}{4} \right) \right]$$

$$= \pi(e^2 - 1) - \pi \left( e^2 - \frac{e^2}{2} + \frac{1}{2} \right)$$

$$= \frac{\pi}{2}(e^2 - 3)$$

$$\begin{aligned} \text{c) i) } f(x) &= e^{-x} \left( 1 + x + \frac{x^2}{2!} + \dots + \frac{x^n}{n!} \right) \\ \therefore f'(x) &= e^{-x} \left( 1 + \frac{2x}{2!} + \frac{3x^2}{3!} + \dots + \frac{nx^{n-1}}{n!} \right) \\ &\quad - e^{-x} \left( 1 + x + \frac{x^2}{2!} + \dots + \frac{x^n}{n!} \right) \end{aligned}$$

$$\therefore f'(x) = e^{-x} \left( 1 + x + \frac{x^2}{2!} + \dots + \frac{x^{n-1}}{(n-1)!} \right)$$

$$- e^{-x} \left( 1 + x + \frac{x^2}{2!} + \dots + \frac{x^n}{n!} \right)$$

$$= -e^{-x} \left( \frac{x^n}{n!} \right) < 0 \text{ for all } x \geq 0.$$

$$\text{since } e^{-x} > 0 \text{ and } \frac{x^n}{n!} > 0.$$

hence the curve is decreasing for all  $x \geq 0$  as  $f'(x) < 0$ .

ii) When  $x=0$   $f(0) = 1$

and since the curve is always decreasing  $f(x) \leq 1$ .

$$\therefore e^{-x} \left( 1 + x + \frac{x^2}{2!} + \dots + \frac{x^n}{n!} \right) \leq 1$$

$$\therefore 1 + x + \frac{x^2}{2!} + \dots + \frac{x^n}{n!} \leq \frac{1}{e^{-x}}$$

$$\text{hence } e^x \geq 1 + x + \frac{x^2}{2!} + \dots + \frac{x^n}{n!}.$$

### Question 16.

$$\text{a) } \int \frac{e^{2x}}{e^x - 1} dx = \int \frac{e^x \cdot e^x}{e^x - 1} dx$$

$$\text{Let } u = e^x - 1 \rightarrow du = e^x dx$$

$$\therefore I = \int \frac{u+1}{u} du = \int 1 + \frac{1}{u} du$$

$$= u + \ln u$$

$$= e^x - 1 + \ln(e^x - 1) + C_1$$

$$= e^x + \ln(e^x - 1) + C_2$$

**Question 16 continued:**

b) i)  $xy = c^2 \rightarrow y = c^2 x^{-1}$

$\therefore y' = \frac{c^2}{x^2} \rightarrow m = -\frac{1}{p^2}$  when  $x = cp$

Equation of tangent at  $P$  :

$y - \frac{c}{p} = -\frac{1}{p^2}(x - cp)$

$p^2 y - cp = -x + cp$

$\therefore x + p^2 y = 2cp$

ii) since  $OT$  is perpendicular tangent at  $P$  it has  $p^2$  as its gradient.

$\therefore$  equation of  $OT$  is  $y = p^2 x$ , and since  $T(x_1, y_1)$

lies on  $OT$  then  $y_1 = p^2 x_1$ .

iii) Since  $T$  satisfies the equation of the tangent:

$x_1 + p^2 y_1 = 2cp$

and from (ii)  $p^2 = \frac{y_1}{x_1}$

$\therefore x_1 + \frac{y_1}{x_1} \cdot y_1 = 2c \sqrt{\frac{y_1}{x_1}}$

$\therefore x_1^2 + y_1^2 = 2c x_1 \sqrt{\frac{y_1}{x_1}}$

$\therefore x_1^2 + y_1^2 = 2c \sqrt{x_1 y_1}$

$\therefore (x_1^2 + y_1^2)^2 = 4c^2 x_1 y_1$

hence the locus of  $T$  is given by

$(x^2 + y^2)^2 = 4c^2 xy$ .

c) i)  $I_n = \int_0^1 x^{2n+1} e^{-x^2} dx = \int_0^1 x^{2n} \cdot x e^{-x^2} dx$

$u = x^{2n}, \quad v' = x e^{-x^2}$

$u' = 2n x^{2n-1}, \quad v = -\frac{1}{2} e^{-x^2}$

$\therefore I_n = \left[ -\frac{1}{2} x^{2n} e^{-x^2} \right]_0^1 + n \int_0^1 x^{2n-1} e^{-x^2} dx$

$= -\frac{1}{2} e^{-1} - 0 + n I_{n-1}$

$\therefore I_n = -\frac{1}{2e} + n I_{n-1}$  -----\*

ii)  $I_0 = \int_0^1 x e^{-x^2} dx$

$= \left[ -\frac{1}{2} e^{-x^2} \right]_0^1 = -\frac{1}{2} (e^{-1} - e^0)$

$= -\frac{1}{2} \left( \frac{1}{e} - 1 \right) = \frac{1}{2} - \frac{1}{2e}$

also  $I_1 = -\frac{1}{2e} + I_0$

$= -\frac{1}{2e} + \frac{1}{2} - \frac{1}{2e} = \frac{1}{2} - \frac{1}{e}$

iii) When  $n=1$ :  $LHS = 1 + \frac{1}{1!} = 2$

$RHS = e - \frac{2e I_1}{1!}$

$= e - 2e \left( \frac{1}{2} - \frac{1}{e} \right)$

$= e - e + 2 = 2 = LHS$

hence true for  $n=1$

Assume true for  $n=k$  :

$1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \dots + \frac{1}{k!} = e - \frac{2e I_k}{k!}$

Prove true for  $n=k+1$  :

i.e.  $S_k + T_{k+1} = S_{k+1}$  (where  $S_{k+1} = e - \frac{2e I_{k+1}}{(k+1)!}$ )

$LHS = e - \frac{2e I_k}{k!} + \frac{1}{(k+1)!}$

$= e - \frac{2e(k+1) I_k}{(k+1)!} + \frac{1}{(k+1)!}$

$= e + \frac{1 - 2e(k+1) I_k}{(k+1)!}$

$$= e + \frac{1 - 2e \left( \frac{1}{2e} + I_{k+1} \right)}{(k+1)!}$$

$$= e + \frac{1 - 1 - 2eI_{k+1}}{(k+1)!}$$

$$= e - \frac{2eI_{k+1}}{(k+1)!} = S_{k+1}$$

Hence by induction is true for all  $n \geq 1$ .

**NOTE:**  $I_{k+1} = -\frac{1}{2e} + (k+1)I_k$  using  $\square$ \*

$$\therefore (k+1)I_k = \frac{1}{2e} + I_{k+1}$$

iv)  $0 \leq I_n \leq 1$ , hence as  $n \rightarrow \infty$ ,  $\frac{I_n}{n!} \rightarrow 0$

$$\therefore 1 + \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \dots + \frac{1}{n!} = e - 2e \times 0 = e$$

$$\therefore \frac{1}{1!} + \frac{1}{2!} + \frac{1}{3!} + \dots = e - 1$$